

SHORT COMMUNICATION

Detection of *Salmonella* sp. and *Escherichia coli* on Chicken Meat at Tamiang Layang Market

AKHMAD RIZALDI¹ AND ENGKI ZELPINA^{2*}

¹*Veterinary Medical Expert of the Department of Agriculture in East Barito Regency, Central Agricultural State Polytechnic of Payakumbuh,*
²*Jl. Raya Negara, KM 7 Tanjung Pati, Harau, Fifty Cities, 26271*

Salmonella sp. and *Escherichia coli* (*E. coli*) are the two most important pathogens because they are indicators of food safety and sanitation indicators, because they can potentially pose a high risk of foodborne disease. This study aims to look at the prevalence of *Salmonella* sp. and *E. coli* in the Tamiang Layang Market as a supervision of food safety. A total of 6 chicken breast samples were taken at all chicken traders in the Tamiang Layang market. Testing the presence of *Salmonella* sp. and *E. coli* using MC-Media Pad. The existence of *Salmonella* sp. and *E. coli* in chicken meat at the Tamiang Layang market were 66.6% and 83.3%. The need to improve hygiene and sanitation for chicken traders.

Key words: chicken meat, *Escherichia coli*, market, *Salmonella* sp.

Salmonella sp. dan *Escherichia coli* (*E. coli*) merupakan dua patogen terpenting karena merupakan indikator keamanan pangan dan indikator sanitasi pada pangan asal hewan, karena berpotensi menimbulkan risiko tinggi penyakit bawaan makanan pada pangan asal hewan. Penelitian ini bertujuan untuk melihat prevalensi *Salmonella* sp. dan *E. coli* di Pasar Tamiang Layang sebagai pengawasan keamanan pangan asal hewan. Sebanyak 6 sampel dada daging ayam yang diambil pada semua pedagang ayam di pasar Tamiang Layang. Pengujian sampel tentang keberadaan *Salmonella* sp. dan *E. coli* menggunakan MC-Media Pad. Keberadaan *Salmonella* sp. dan *E. coli* pada daging ayam di pasar Tamiang Layang masing-masing sebesar 66,6% dan 83,3%, Perlunya meningkatkan higienitas dan sanitasi para pedagang ayam.

Kata kunci: daging ayam, *Escherichia coli*, pasar, *Salmonella* sp.

Food is the most important basic human need in maintaining a healthy body, growth, and improving people's intelligence. Meat is an important food ingredient in meeting nutritional needs. Chicken meat is one of the sources of high-quality animal protein that is much in demand by the community because it is easily digested, can be accepted by the majority of people and the price is relatively cheap.

Chicken meat is muscle tissue obtained from chicken which is commonly used for food consumption purposes. The speed at which meat breaks down depends on the number of initial microbes. The greater the number of initial microbes in the meat, the faster the damage will be. Microbial contamination in foodstuffs of animal origin and its products is a problem of major concern for consumers. There are many potential critical points for the occurrence of contact and entry of microbes into food from animal origin and their processed products,

therefore it is necessary to identify microbial contaminants in maintaining food safety from farm to dining table (safe from farm to table).

Contamination of pathogenic microbes in animal foods such as chicken is a health problem that needs attention. Several types of pathogenic bacteria that can contaminate chicken meat are *Salmonella* sp. and *Escherichia coli* (*E. coli*). Both types of bacteria are the main cause of Foodborne disease. Foodborne disease is a disease that occurs due to consuming food contaminated with microbes such as bacteria with common symptoms of abdominal pain, vomiting, diarrhea, dizziness, convulsions, and fever (CDC 2018). Food safety assurance or foodstuffs have become demands along with the increasing public awareness of health.

Tamiang Layang Traditional Market is the main market that provides chicken meat every day for the people of East Barito. At the Tamiang Layang traditional market, there are eight stalls selling chicken meat. Chicken meat can be contaminated in the slaughterhouse, at the point of sale, and during

*Corresponding author: Phone: +62-; Fax: -; Email: engkizelpina03@gmail.com

Table 1 Presence of *Salmonella* sp. and *Escherichia coli* in chicken

Sample	<i>Salmonella</i> sp.	<i>Escherichia coli</i>
Sample 1	+	+
Sample 2	-	+
Sample 3	+	+
Sample 4	-	-
Sample 5	+	+
Sample 6	+	+

Table 2 Prevalence of *Salmonella* sp. and *Escherichia coli* in chicken

Bacteria	Prevalence (%) (n=6)	
	Positive	Negative
<i>Salmonella</i> sp.	4 (66.6)	2 (33.4)
<i>Escherichia coli</i>	5 (83.3)	1 (16.7)

processing with incomplete cooking.

The possibility of *Salmonella* sp. and *E. coli* in chicken meat during the sales process at the Traditional Tamiang Layang market, it is necessary to conduct research to detect the presence or absence of *Salmonella* sp. bacterial contamination and *E. coli* in chicken sold in the Tamiang Layang traditional market. Information about the contamination of these two bacteria in chicken meat products sold at the Tamiang Layang traditional market will be able to increase the awareness of the Tamiang Layang community in particular and East Barito in general in buying and consuming chicken meat sold in traditional markets in East Barito.

A total of 6 chicken meat samples were obtained from chicken traders at the Tamiang Layang market, East Barito. The samples taken were put into a sterile plastic and stored in a cooler box containing an ice pack and immediately taken to the laboratory for bacteriological testing.

Isolation and Presence of *Escherichia coli*. A total of 50 grams of chicken thigh meat was weighed then added with 450 mL of Phosphate Buffered Saline (FBS), after which it was homogenized using a stomacher. Prepare the MC-Media Pad *E. coli* test kit by opening the aluminum cap, then opening the transparent pad cover and then dropping 1 mL of a mixture of chicken meat with homogeneous FBS earlier. Enter it slowly diagonally, then close it again. Mc-Media Pad *E. coli* was inserted into a petri dish, given a code, then incubated at $35 \pm 1^\circ \text{C}$ for 24 ± 2 hours. AOAC OMA 966.24 (Cer 070901) was used as a positive control. Colonies and the presence of *E. coli* will appear as purplish-red colonies (Millipore 2017).

Isolation of the presence of *Salmonella* sp. A total of 10 grams of thigh meat samples were weighed then

mixed with 90 mL of Phosphate Buffered Saline (FBS) solution, after which it was homogenized with a stomacher. Prepare the MC-Media Pad *Salmonella* test kit by opening the aluminum cover, then opening the transparent pad cover and then dropping 1 mL of the chicken meat mixture with homogeneous FBS earlier. Insert it slowly diagonally, then close it again. Mc-Media Pad *Salmonella* sp. put into a petri dish, given a code, then incubated at $35 \pm 1^\circ \text{C}$ for 24 ± 2 hours. The colony and the existence of *Salmonella* sp. will appear as a light blue colony (Millipore 2017).

Based on bacteriological analysis in the laboratory to determine the presence of *Salmonella* and *E. coli* from chicken meat obtained from the Tamiang Layang market, 6 samples were examined with the following results in Table 1.

Table 1 shows that the presence of *Salmonella* sp. and *E. coli* were positive in 4 samples, i.e. sample (1,3,5,6), whereas in sample 2 it was negative *Salmonella* sp. and positive in *E. coli* and in sample 4 negative for both. *Salmonella* sp. and *E. coli* are food pathogenic bacteria that are transmitted by fecal or oral. The presence of these bacteria in food products of animal origin is caused by many factors of livestock/animal origin, the process of slaughtering, the process of transportation of the product to the market/retail, as well as handling at the sales location (Nel *et al.*, 2004; Regalado-Pineda *et al.*, 2020) Based on studies conducted on the wet market environment and small-scale processing plants in Perlis and Penang, Malaysia, *Salmonella* sp. not only in meat but also found in utensils and equipment used in processing, washing containers, soaking containers, tables, drains and floors around wet markets and processing plants. This suggests that this source is a potential risk for transmission of *Salmonella* sp. on chicken meat and its

environment (Nidaulah *et al.* 2017). Poor sanitation and hygiene conditions and high humidity in the slaughterhouse and chicken meat processing industry are ideal for biofilm formation by *Salmonella* sp. Bacteria. These biofilms are long-lasting and tend to protect *Salmonella* sp. from cleaners, so it is very risky to cross-contaminate *Salmonella* sp. in chicken and chicken meat shops (Smith *et al.* 2007).

Based on Table 2, the prevalence of *Salmonella* sp. 4 (66%) and *E. coli* 5 (83,35). The prevalence of *E. coli* is higher than that of *Salmonella* sp. high prevalence *E. coli* in chicken meat shows the low application of sanitation and personal hygiene in producing chicken meat at the Tamiang Layang Market. *Salmonella* sp. and *E. coli* are indicators of food safety and sanitation indicators on the food of animal origin. Meat is one of the main sources of contamination from *Salmonella* sp. and *E. coli*, therefore the importance of controlling and surveillance of the presence of these pathogens in food of animal origin (Hedican *et al.* 2007; Kirk *et al.* 2015).

The condition of the Tamiyang Layang market, which is located in an open stall for sales of chicken meat, is carried out on the sale site until the carcass is ready for sale without refrigeration resulting in cross-contamination of the presence of *Salmonella* sp. and *E. coli*. Sales of chicken meat in open markets without refrigeration have a chance to be contaminated by *Salmonella* sp. and *E. coli* is higher than sales in closed and refrigerated spaces (Nidaulah *et al.* 2016). In addition, the presence of *Salmonella* sp. and *E. coli* are also reported to be found in several processed chicken meat such as grilled chicken, chicken sausage and shredded chicken in chicken porridge (Amiruddin *et al.* 2017; Kartika *et al.* 2014; Zelpina *et al.* 2018).

Poor sanitary conditions in the market for sales in tropical countries also have the potential for high risk of contamination by *Salmonella* sp. and *E. coli* which can cause foodborne disease and is a zoonotic disease for the community. The presence of *Salmonella* sp. and *Escherichia coli* in chicken meat, not only showed poor sanitary conditions during slaughter but also showed the health status of poultry as carrier carriers against *Salmonella* sp. and *Escherichia coli*.

The presence of *Salmonella* sp. and *Escherichia coli* in chicken meat sold at the Tamiang Layang Market, stated that the implementation of sanitary and personal hygiene in every stage of the chicken meat production process has the potential to cause disease in the community because it is a foodborne disease.

REFERENCES

- Amiruddin RR, Darniati, Ismail. 2017. Isolation and Identification of *Salmonella* sp. in Roasted Chicken from Restaurant in Syiah Kuala, Banda Aceh. JIMVET. 1(3):265-274. doi: 10.21157/jim%20vet.v1i3.3152.
- [CDC] Centers for Disease Control and Prevention. 2018. *Foodborne Disease* [internet]. [diunduh 2020 July 6]. Tersedia pada: <https://www.cdc.gov/foodsafety/foodborne-germs.html>.
- Crump JA, Medalla FM, Joyce KW, Krueger AL, Hoe kstra RM, Whichard JM, Barzilay EJ. 2011. Antimicrobial resistance among invasive nontyphoidal *Salmonella* enterica isolates in the United States: National Antimicrobial Resistance Monitoring System, 1996 to 2007. *Antimicrob Agents Chemother.* 55:1148-1154. doi: 10.1128/AAC.01333-10.
- Doaa MAEA. 2013. Detection of *Salmonella typhimurium* in retail chicken meat and chicken giblets. *Asian Pac J Trop Biomed.* 3(9):678–681. doi: 10.1016/S2221-1691(13)60138-0.
- Furukawa I, Ishihara T, Teranishi H, Saito S, Yatsuyanagi J, Wada E, Kumagai Y, Takahashi S, Konno T, Kashio H, Kobayashi A, Kato N, Hayashi KI, Fukushima K, Ishikawa K, Horikawa K, Oishi A, Izumiya H, Ohnishi T, Konishi Y, Kuroki T. 2017. Prevalence and Characteristics of *Salmonella* and *Campylobacter* in Retail Poultry Meat in Japan. *Jpn J Infect Dis.* 70(3):239-247. doi: 10.7883/yoken.JJID.2016.164.
- Hedican E, Smith K, Jawahir S, Scheftel J, Kruger K, Birk R, *et al.* 2009. Multistate outbreaks of *Salmonella* infections associated with live poultry – United States, 2007. *Morb Mortal Wkly Rep.* 58:25–34.
- Kartika E, Khotimah S, Yanti AH. 2014. Detection of Food Safety Indicator Bacteria in Sausages Chicken Meat at Pontianak's Flamboyant Market. *Protobiont.* 3(2): 111-119.
- Khan AS, Georges K, Rahaman S, Abdela W, Adesiyun AA. 2018. Prevalence and serotypes of *Salmonella* spp. on chickens sold at retail outlets in Trinidad. *PLoS ONE.* 13(8): e0202108. doi: 10.1371/journal.pone.0202108.
- Kirk MD, Pires SM, Black RE, Caipo M, Crump JA, Devleeschauwer B, *et al.* 2015. World health organization estimates of the global and regional disease burden of 22 foodborne bacterial, protozoal, and viral diseases, 2010: a data synthesis. *PLoS Med.* 12(12):e1001921. doi: 10.1371/journal.pmed.1001921.
- Lee MB, Middleton D. 2003. Enteric Illness in Ontario, Canada, from 1997 to 2001. *J Food Prot.* 66:953-996. doi: 10.4315/0362-028x-66.6.953.
- Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M,

- O'Brien SJ, Jones TF, *et al.* 2010. The global burden of non-typhoidal *Salmonella* gastroenteritis. *Clin Infect Dis.* 50:882-889. doi: 10.1086/650733.
- Millipord. 2017. Instruction Manual. MC-Media Pad E. Coli and Coliform. Germany.
- Moawad AA, Hotzel H, Awad O, Tomaso H, Neubauer H, Hafez HM, El-Adawy H. 2017. Occurrence of *Salmonella enterica* and *Escherichia coli* in raw chicken and beef meat in northern Egypt and dissemination of their antibiotic resistance markers. *Gut pathogens.* 9:57. doi: 10.1186/s13099-017-0206-9.
- Nel S, Lues JFR, Buys EM, Venter P. 2004. Bacterial populations associated with meat from the deboning room of a high throughput red meat abattoir. *Meat Sci.* 66:667-674. doi: 10.1016/S0309-1740(03)00187-6.
- Nidaulah H, Mohd OAK, Ahmad R, Nurul H, dan Saima S. 2016. Analysis of *Salmonella* Contamination in Poultry Meat at Various Retailing, Different Storage Temperatures and Carcass Cuts. *International Journal of Poultry Science.* 15(3):111-120. doi: 10.3923/ijps.2016.111.120.
- Nidaullah H, Abirami N, Shamila-Syuhada AK, Chuah LO, Nurul H, Tan TP, Abidin F, Rusul G. 2017. Prevalence of *Salmonella* in poultry processing environments in wet markets in Penang and Perlis, Malaysia. *Veterinary world.* 10(3):286-292. doi: 10.14202/vetworld.2017.286-292.
- Regalado-Pineda ID, Rodarte-Medina R, Resendiz-Nava CN, Saenz-Garcia CE, Castañeda-Serrano P, Nava GM. 2020. Three-Year Longitudinal Study: Prevalence of *Salmonella Enterica* in Chicken Meat is Higher in Supermarkets Than Wet Markets from Mexico. *Foods (Basel, Switzerland).* 9(3):264. doi: 10.3390/foods9030264.
- Smith DP, Northcutt JK, Cason JA, A. Hinton JR, Buhr RJ and Ingram KD. 2007. Effect of external or internal faecal contamination on numbers of bacteria on prechilled broiler carcasses. *Poult Sci.* 86:1241-1244. doi: 10.1093/ps/86.6.1241.
- Zelpina E, Purnawarman T, Lukman DW. 2018. Occurance *Salmonella sp.* in Shredded Chicken Meat of Chicken Porridge Sold around Bogor Agricultural University, Dramaga, Bogor. *Jurnal Penelitian Pascapanen Pertanian.* 15(2):73-79. doi: 10.21082/jpasca.v15n2.2018.73-79.